



KAMIYA BIOMEDICAL COMPANY

Alkaline Phosphatase-NH₂ Labeling Kit

**For the rapid Alkaline Phosphatase labeling of protein
for EIA and Immunoblotting/Immunostaining**

Cat. No. KT-220

For Research Use Only.

PRODUCT INFORMATION

Alkaline Phosphatase-NH₂ Labeling Kit Cat. No. KT-220

PRODUCT

Alkaline Phosphatase-NH₂ Labeling Kit is designed for the simple and rapid preparation of alkaline phosphatase-labeled IgG for enzyme immunoassays (EIA) and immunoblotting / immunostaining. It can also be used for the preparation of alkaline phosphatase-labeled antigen for competitive EIA. NH₂-reactive Alkaline Phosphatase (AP) (a component of this kit) has activated ester groups and can easily make a covalent bond with an amino group of the target molecule without any activation process. If the target is a small molecule, the conjugate can be purified with the Filtration Tube included in this kit. The Filtration Tube is also used for removing small molecules such as Tris buffer and amine compounds that may interfere with this assay or labeling reaction. This kit contains all of the necessary reagents for alkaline phosphatase labeling, including the storage buffer for the conjugates.

For research use only.

COMPONENTS

- | | |
|--------------------------------|------------|
| • NH ₂ -reactive AP | 3 X 100 µg |
| • Reaction Buffer | 200 µL |
| • Wash Buffer | 4 mL |
| • Storage Buffer | 4 mL |
| • Filtration Tubes | 3 tubes |

Materials or equipment required but not provided

- 0.5 mL microtubes
- 10 µL and 200 µL adjustable pipettes
- Microcentrifuge
- 37°C Incubator

SAMPLE REQUIREMENT

Proteins: Molecular weight >50,000; amount: 50-200 µg (IgG)

Small Molecule: Molecular weight <5,000

PROCEDURE

Labeling of Sample (IgG)

1. Add 100 µL of Wash Buffer and the sample solution containing 50-200 µg of IgG to the Filtration Tube.
2. Centrifuge at 8,000-10,000 g for 10 minutes. Add 100 µL Wash Buffer and centrifuge again.
3. Add 10 µL Reaction Buffer to the NH₂-reactive AP and dissolve it by pipetting up and down.
4. Transfer the solution containing the NH₂-reactive AP onto the membrane where the IgG is concentrated.
5. Rinse the entire surface of the membrane with the solution by pipetting and incubate the tube at 37°C for 2 hours.
6. Add 190 µL of Storage Buffer and pipette 10 to 15 times to recover the conjugate. Transfer the solution to a 0.5 mL tube and store the solution at 4°C.

Precautions

IgG or alkaline phosphatase-conjugated IgG is always on the filter membrane of the filtration tube during the labeling process. If the IgG solution contains other proteins with molecular weights larger than 10,000, such as BSA or gelatin, purify the IgG solution prior to labeling with this kit. IgG solution can be purified by IgG purifications kits (not included in this kit). If the IgG solution contains small insoluble materials, centrifuge the solution and use the supernatant for labeling.

The recommended amount of IgG is 100 µg in a volume of 100 µL or less. If the antibody concentration is lower than 0.5 mg/mL, repeat step 1 and 2 until the total IgG accumulation becomes 50-200 µg. If the volume of the filtrate

becomes 400 μL or more during the accumulation process, discard the filtrate prior to going on to the next centrifuge step.

If solution still remains on the membrane after centrifugation, spin another 5 minutes or increase the centrifuge speed.

The concentration of the conjugate is 0.5-1.3 mg/mL. Dilute the alkaline phosphatase-labeled IgG to prepare a solution with an appropriate concentration prior to using it for enzyme immunoassay, immunoblotting or immunostaining. One to three molecules of alkaline phosphatase should be introduced onto one IgG molecule. Unconjugated alkaline phosphatase should not interfere with normal immunoassays. If purification is necessary, use a gel permeation column or an affinity column for IgG.

Generally the alkaline phosphatase-labeled IgG in Storage Buffer is stable for at least 2 months at 4°C. For longer storage, store at -20°C. However, it is important to note that the stability will depend on the sample itself.

Labeling of Small Molecule

1. Prepare 50 μL of a 1 mM amine compound solution with Reaction Buffer and add the solution to a tube of NH_2 -reactive AP. Pipette several times to mix and incubate at 37°C for 1 hour.
2. Add 100 μL Wash Buffer to the reaction solution and transfer the entire solution to a Filtration Tube.
3. Centrifuge at 8,000-10,000 g for 10 minutes. Discard the filtrate and add 200 μL of Wash Buffer to the tube, and centrifuge at 8,000-10,000 g for 10 min. Add 200 μL Wash buffer and centrifuge again.
4. Add 200 μL of Storage Buffer and pipette 10-15 times to recover the conjugate. Transfer the solution to a 0.5 mL tube and store the solution at 4°C.

Precautions

If the amine compound does not dissolve in aqueous solution, dissolve it with DMSO to prepare a 10 mM solution. Mix 5 μL of the solution with 45 μL Reaction Buffer.

If solution still remains on the membrane after the centrifugation, spin again for 5 minutes or increase the centrifuge speed.

The concentration of the conjugate is 400-500 $\mu\text{g}/\text{mL}$. One to two target molecules should be conjugated with one alkaline phosphatase molecule.

The alkaline phosphatase-labeled small molecule is stable for at least 6 months at 4°C.

STORAGE

Store all components at 4°C. Stable until the expiration date at 4°C with protection from moisture.

FAQ

Q. Can I use this kit for Fab or Fab' labeling?

A. Yes, you can label Fab or Fab' using this kit. The recovery of the conjugate should be over 80%.

Q. Can I use this kit for other proteins or peptides?

A. Yes, if the molecular weight is greater than 50,000 or less than 5,000 and it has a reactive primary or secondary amino group. If the molecular weight is greater than 50,000, follow the labeling protocol for IgG and use 0.5-1 nmol of sample protein. If the molecular weight is less than 5,000, follow the labeling protocol for small molecules.

Q. Can I use this kit on oligonucleotides?

A. Yes, if the molecular weight is less than 5,000 and it has a reactive primary or secondary amino group. Follow the labeling protocol for small molecules.

Q. What is the minimal amount of IgG that can be labeled with this kit?

A. The minimal amount is 50 μg . There is no significant difference in sensitivity and background between 50 μg and 200 μg of IgG. However, even 10 μg of IgG can still be labeled using this kit, the background will be higher.

Q. How many alkaline phosphatase molecules per IgG are introduced?

A. Average number of alkaline phosphatase molecules per IgG is 1 to 3.

Q. Does unconjugated NH₂-reactive AP still have an activated ester after the labeling reaction to IgG?

A. No, it is completely hydrolyzed during the reaction.

Q. Does NH₂-reactive AP form an oligomer during the labeling reaction?

A. No, since all amino groups of NH₂-reactive AP are blocked, no oligomerization is possible.

Q. Do I have to use Storage buffer included in the kit?

A. No, you do not have to use Storage Buffer from the kit. You can choose a buffer that is appropriate for your experiment.

FOR RESEARCH USE ONLY

KAMIYA BIOMEDICAL COMPANY

12779 Gateway Drive, Seattle WA 98168

Tel: (206) 575-8068 Fax: (206) 575-8094

Email: techsupport@kamyabiomedical.com

www.kamyabiomedical.com